



Bacteriological Analysis of Surface Water Obtained from Uke River in Karu, Nasarawa State, Nigeria

J. O. Ihuma¹, L. Y. Adogo^{1*}, T. D. Malgwi¹, J. I. Maikenti¹ and A. Z. Koggie¹

¹Department of Biological Sciences, Faculty of Science and Technology, Bingham University, Karu, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Authors JOI and TDM designed the study, wrote the protocol and wrote the first draft of the manuscript. Author LYA reviewed the experimental design and all drafts of the manuscript. Authors JOI and TDM managed the analyses of the study. Authors JIM and AZK performed the statistical analysis. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2016/30518

Editor(s):

(1) Umesh Gangishetti, Department of Biology, Emory University, Atlanta, USA.

Reviewers:

(1) Fazl Fen, Yüzüncü Yıl University, Turkey.

(2) Naeem Khan, Kohat University of Science and Technology, Pakistan.

(3) Mahmoud M. M. Zaky, Port Said University, Egypt.

Complete Peer review History: <http://prh.sdiarticle3.com/review-history/17348>

Original Research Article

Received 14th November 2016
Accepted 19th December 2016
Published 26th December 2016

ABSTRACT

Aim: To determine the bacteriological properties of the surface water obtained from Uke River.

Place and Duration of Study: This study was carried out in Uke Community, Karu LGA, Nasarawa state and the Department of Biological sciences, Bingham University, Karu, between March and April 2016.

Materials and Methods: Water samples were collected randomly from five points at the Uke River for five weeks. The total heterotrophic bacteria count, fecal coliform count and total coliform count were used to determine bacterial contamination. Biochemical tests and gram reaction was used to identify the bacterial isolates.

Results: Six bacterial genera which include *Salmonella* spp, *Staphylococcus* spp, *Escherichia* spp, *Pseudomonas* spp, *Enterococcus* spp and *Klebsiella* spp were isolated from the water samples. *Escherichia coli* had the highest frequency of 23(43%). The mean total heterotrophic count, total

*Corresponding author: E-mail: adogolillian@gmail.com;

coliform count and fecal coliform count were 2.3×10^2 , 2.4×10^2 , and 2.6×10^2 respectively.

Conclusion: The bacteriological analysis of the surface water indicates the presence of bacteria which suggests the water is not fit for consumption without proper processing.

Keywords: Bacteria; Uke River; fecal coliform; water.

1. INTRODUCTION

The accessibility to safe water and sanitary means of excreta disposal are universal needs and indeed basic human rights. However, many of the world's population lack access to adequate and safe water [1].

Water is vital to the existence of all living organisms; however, this valued resource is increasingly being threatened as a result of upsurge in human population. Consequently, the demand for high quality water for the both domestic and economic purposes has been noticeably on constant increase [2]. Water quality is defined in terms of its chemical, physical and biological contents. The water quality of rivers changes with seasons, human activities and geographical areas. These factors provide basic scientific information about water quality parameters, ecological relevance and toxicological threshold values to protect specific water user [3].

Rivers are the source of freshwater to man. Flooding remains a source of pollution for rivers in Nigeria. Surface waters are heavily polluted from several sources especially after rainfall [4]. Sources of pollution include runoff water from farmlands carrying fertilizers, manure, animal and human waste matter, motor oils from the highways and lots of trash from gutters and drainages. World Health Organization (WHO) essential parameters of drinking water quality are fecal *Escherichia coli* and total coliforms, chlorine residue, turbidity, pH, dissolved oxygen content, and temperature [5].

According to UNICEF report, about 800 million people in Asia and Africa are living without access to safe drinking water. Consequently this has caused many people to suffer from various diseases [6]. However, access to safe drinking water has improved over the last decades in almost every part of the world especially Nigeria, but approximately 1.1 billion people still lack access to safe water and over 2.6 billion worldwide lack access to adequate water and sanitation which causes water illnesses such as

Cholera, diarrheal disease, Botulism, *E. coli* infection, Dysentery, Legionellosis, Leptospirosis, Salmonellosis, Typhoid fever, and Vibrio illness [7].

The greatest risk to public health from microbes in water is associated with consumption of drinking-water that is contaminated with human and animal excreta [8]. Human faeces can contain a variety of intestinal pathogens which cause diseases ranging from mild gastro-enteritis to the serious dysentery, cholera and typhoid. The most predominant waterborne disease, diarrhea, has an estimated annual incidence of 4.6 billion episodes and causes 2.2 million deaths every year [9].

Faecal coliform bacteria are used as an indicator for the presence of any of these water borne pathogens. The presence of these indicative organisms is evidence that the water has been polluted with faeces of humans or other warm-blooded animals [4,10]. The Uke River is a source of drinking water to the members of the community and it's characterized with human activities such as bathing, washing, swimming and construction of blocks. Although some studies have been carried out on surface water resources in Nigeria, there is paucity of information on the bacteriological analysis of Uke River which informed this study.

2. MATERIALS AND METHODS

2.1 Study Area

This study was carried out in Uke local government area, Nasarawa State. This community is located in the middle belt of Nigeria at longitude $8^{\circ}32'N$ $8^{\circ}18'E$ and Latitude $8.533^{\circ}N$ $8.300^{\circ}E$ and is characterized by a tropical sub-humid climate with two distinct seasons; wet and dry seasons. Monthly temperature ranges from $20^{\circ}C$ to $34^{\circ}C$ and annual rainfall ranges from 1100 mm to about 2000 mm. Uke is located about 26 kilometers South-East of Abuja, the country's capital town. It is regarded as a suburb of Abuja due to its close proximity to Nigeria's Federal Capital Territory [11].




Fig. 1. Map of Nasarawa State highlighting the study area (Uke) in Karu Local Government
 Source: Chindo et al, 2014

2.2 Sample Collection

Water samples were collected from five (5) different water sites indicated as location A, B, C, D, E and a COMPOSITE (F) which includes all samples. The water samples were collected at these sites because the inhabitants believe that the cleanest water can be obtained from these points. Water samples from these locations were collected in duplicates into sterile glass bottles (250 ml) which were labeled appropriately and transported to the laboratory for bacteriological analysis.

2.3 Bacteriological Analysis

Blood agar and MacConkey agar were prepared using manufacturers direction [12]. Bacteria isolates were characterized on the basis of the colonial morphology and Gram stain reaction. Biochemical tests such as catalase, Coagulase, Motility, Indole, and Oxidase tests were carried out [13].

2.3.1 Total heterotrophic bacteria count

The spread plate method was used. Ten-fold serial dilution of each water sample was prepared aseptically in physiological saline of 10^{-1} up to 10^{-4} and 0.1 ml aliquot of each dilution was plated on Nutrient agar plates in triplicate. All incubations were conducted at 37°C for 24

hrs under aerobic conditions and plates containing 30 to 300 colonies were selected and counted. The number of colony-forming units per ml (cfu/ml) was calculated by multiplying the number of colonies by the dilution factor. Also, sub-culture was carried on MacConkey agar and Mannitol Salt agar for identification of bacteria species.

2.3.2 Total coliform count

This was determined by Most Probable Number (MPN) index technique using the three tube assay (3-3-3 regimen). Ten-fold serial dilution of 10^{-1} to 10^{-5} was prepared. The first set of five tubes had 10 ml of double strength broth (MacConkey broth), the second and third set had 10ml single strength broth (Lactose broth). All the tubes contained Durham tubes. The three set of tubes received 10 ml, 1 ml and 0.1 ml of water samples. They were carefully labeled and incubated at 37°C for 24 hours for estimation of total coliform. Acid production was determined by color change in tubes from reddish purple to yellow and gas production was checked for by entrapment of gas in the durham tubes [13].

2.3.3 Faecal coliform count

Faecal coliform count was determined using Eosin Methylene Blue medium employing the streaking culture technique. A loopful of broth

from positive tubes was streaked onto EMB agar plate for pure cultures. The plates were incubated at 37°C for 24 hrs. Colonies on EMB agar plate were further identified as fecal coliforms. On Eosin Methylene Blue (EMB) agar, *E. coli* strains appeared as greenish metallic sheen colonies [14].

3. RESULTS

Table 1 shows the morphological, biochemical characteristics and Gram reaction of the bacterial isolates.

Table 2 represents the frequency of bacteria isolates recorded in Uke River. *Escherichia coli* had the highest frequency of 23(43%). *Pseudomonas* spp, *Staphylococcus aureus* and *Klebsiella* spp had a frequency of 18%, 16 %, and 12% respectively. *Streptococcus* spp and *Bacillus* spp had a prevalence of 6% and 4% respectively. *Enterococcus* spp and *Salmonella* spp showed the least occurrence of 2%.

The mean total heterotrophic count was 2.3×10^2 cfu/mL. The mean total coliform count was 2.4×10^2 cfu/100mL. The mean fecal coliform count was 2.6×10^2 cfu/ml as shown on Table 3.

4. DISCUSSION

The bacteriological assessment of Uke River reveals the presence of bacterial contaminants and this is in agreement with the findings of Doughari et al. [15] and May et al. [16] who conducted similar studies in Gudu stream Abuja, and Ebutte River in Ehor community, Edo state, Nigeria.

Total heterotrophic counts ranged from 1.8×10^2 to 2.67×10^2 from the five sampling points of Uke river and these results exceeds the World Health Organization standard for heterotrophic bacteria

in potable which states that the total heterotrophic bacteria count should not be more than 100 cfu/ml [17].

The alarming high number of total coliforms and faecal coliforms per 100 mL obtained from the water samples indicates high level of faecal contamination of the river water which potentially poses a high health risk for the inhabitants of the community. This agrees with Chessbrough [18] who stated that high coliform counts are an indication of high faecal contamination. Faecal streptococci counts indicate more contamination with human excrement than animal excrement.

None of the sampling points of the water sources complied with World Health Organization standard for coliform in water. The total coliform and fecal count for all samples were extremely higher than the WHO standard for coliform bacteria in water which is zero total coliform per 100 ml of water [19].

Similarly, the health guidelines for the use of wastewater in agriculture and aquaculture states that water to be used for irrigation of crops that is likely eaten uncooked and water to be used for sports and public parks in unrestricted regions should not exceed 10³ per 100 mL faecal coliforms [20]. Additionally, the water quality was unacceptable as per EPA's (Environmental Protection Agency) standard of zero (0) fecal colony/100 mL of water to be used in irrigation of any food crops not commercially processed including crops eaten raw [21].

Escherichia coli had the highest frequency (43%) of bacteria isolated from the water which is contrary to the WHO guideline value of zero (0) *E. coli* per 100 mL of drinking water [18]. Similarly, Khalid et al. [21] also recorded a high frequency of *Escherichia coli* in Tigris River, Baghdad.

Table 1. Characterization and identification of isolates from Uke River

Isolates	Cat	Coa	Oxi	Ind	Mot	Gram staining	Morphology	Organism present
1	+	-	-	+	+	-rod	Milky colonies	<i>Escherichia coli</i>
2	+	+	-	-	-	+cocci	Yellow/gold	<i>Staphylococcus</i> spp
3	-	-	-	-	+	-cocci	Milky	<i>Enterococcus</i> spp
4	+	+	-	-	-	-rod	Pink mucoid	<i>Klebsiella</i> spp
5	+	-	+	-	+	-rod	Creamy colonies	<i>Pseudomonas</i> spp
6	+	-	-	-	+	-rod	White colonies	<i>Salmonella</i> spp
7	-	-	-	-	-	+cocci	Milky	<i>Streptococcus</i> spp
8	+	-	-	+	+	+rod	Pale pink	<i>Bacillus</i> spp

Keys: Cat= Catalase, Coa= Coagulase, Oxi= Oxidase, Ind= Indole, Mot= Motility

Table 2. Frequency of bacteria isolated from Uke River

Bacteria	Frequency (%)
<i>Escherichia coli</i>	23(43%)
<i>Staphylococcus aureus</i>	9(16 %)
<i>Enterococcus</i> spp	1(2%)
<i>Klebsiella</i> spp	7(12%)
<i>Pseudomonas</i> spp	10(18%)
<i>Salmonella typhi</i>	1(2%)
<i>Streptococcus</i> spp	3(6%)
<i>Bacillus</i> spp	2(4%)
Total	56

Table 3. Mean total heterotrophic, total coliform and fecal coliform counts of the water samples taken for five weeks from the Uke River

Sampling period	Total heterotrophic count (TH) (cfu/mL)	Total coliform count (TC) (cfu/100mL)	Fecal coliform count ($\times 10^2$ cfu/ml)
Week 1	2.3×10^2	2.9×10^2	2.6×10^2
Week 2	2.5×10^2	2.5×10^2	2.2×10^2
Week 3	2.27×10^2	2.2×10^2	1.2×10^2
Week 4	1.8×10^2	1.69×10^2	2.0×10^2
Week 5	2.67×10^2	2.9×10^2	2.3×10^2
Mean	2.3×10^2	2.4×10^2	2.6×10^2

5. CONCLUSION

The bacteriological analysis of Uke River from this study is a definitive proof that the water is unsafe for consumption as coliform counts were more than the international permissible levels recommended by World Health Organization. Thus, this water is unfit for drinking, bathing, washing of farm produce or for any other agricultural purpose except if it is adequately treated. Public awareness and a total halt of indiscriminate dumping of refuse and feces into the river will go a long way to reduce bacterial contamination.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Dagne T, Assefa D, Aberra G, Woldemariam G, Solomon F, Oliver S. Rapid assessment drinking water quality in the Federal Democratic Republic of Ethiopia. Country Report of Pilot Project Implementation in 2004–2005. Geneva, New York; 2010.
- United Nations Environment Programme. Global Environment Monitoring System/ Water Programme. Water Quality for Ecosystem and Human Health. National Water Research Institute; Burlington, ON, Canada; 2000. ISBN 92-95039-14-9
- Lawson EO. Physico-chemical parameters and heavy metal contents of water from the mangrove swamps of Logos, Lagoon, Nigeria. *Advances in Biological Research*. 2011;5(1):8-21.
- Chukwurah EI. *Aquatic Microbiology*. Otopa Press Limited, Onitsha, Nigeria; 2001.
- World Health Organization. Guidelines for drinking water quality, World Health Organization, Geneva, Switzerland; 2003.
- Tanwir F, Saboor A, Shan MH. Water contamination, health hazards and public awareness-a care of the urban Punjab, Pakistan. *International Journal of Agriculture and Biology*. 2003;(5):560–562.
- Millennium Development Goals (MDGs) report. Water quality; 2008. (Accessed 2010)
- World Health Organization. Guidelines for drinking-water quality. Fourth edition. Geneva, Switzerland; 2011.
- World Health Organization. Water for health, WHO Guidelines for Drinking-water Quality. Geneva, Switzerland; 2010.
- Okpokwasili GB, Akojubi TC. Bacteriological indicators of tropical water quality. *Environmental Toxicology and Water Quality*. 1996;(11):77-82.

11. Available:<http://news1.africanseer.com/templates/default.aspx?a=5202&template=print-article.htm>
12. Cheesbrough M. District laboratory practice for tropical countries, Part 2. Cambridge University Press, Cambridge. 2000;143-154.
13. Fawole MO, Oso BA. Laboratory manual of microbiology. Ibadan, Nigeria (Spectrum Books). MD, USA. Official Method of Water Analysis. 2001;15-45.
14. Burnett SL. Beuchat LR. Human pathogens associated with raw produce and unpasteurized juices, and difficulties in contamination. Journal of Industrial Microbiology and Biotechnology. 2001; 27(6):104-110.
15. Doughari J, Dodo J, Mbuh F. Impact of effluent from Gudu district sewage treatment plant on Gudu stream in Abuja, Nigeria. J. Appl. Sci. Environ. Manage. 2007;11(1):79–83.
16. May NO, Omoigberale JO. Isibor, JI. Izegaegbe MI. Seasonal variation in the bacteriological quality of Ebutte river in Ehor community EDO state, Nigeria. American Journal of Research Communication. 2013;1(7):59-69. ISSN: 2325-4076.
17. World Health Organization. Guidelines for drinking water quality, first addendum to third edition. World Health Organization Press, Geneva, Switzerland. 2006;1.
18. World Health Organization. Guideline of drinking water quality. Health Criteria and Other Supporting Information. WHO, Geneva, Switzerland. 1993;162.
19. World Health Organization. Health guidelines for the use of wastewater in agriculture and aquaculture. Report of a World Health Organization Press. Geneva, Switzerland; 1989.
20. EPA, Guidelines for water reuse. Washington, DC. Environmental Protection Agency; 1992. Technical Report no. EPA/625/R592/004
21. Khalid K, Al-Bayatti KH, Al-Arajy, Al-Nuaemy SH. Bacteriological and physicochemical studies on Tigris river near the water purification stations within baghdad province. Journal of Environmental and Public Health; 2012. Article ID 695253.

© 2016 Ihuma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://prh.sdiarticle3.com/review-history/17348>